

S-asymptotically ω -periodic solutions in the p -th mean for a Stochastic Evolution Equation driven by \mathcal{Q} -Brownian motion

Solym Mawaki Manou-Abi^{*1}, William Dimbour²

¹ *CUFR Mayotte, IMAG-Montpellier, Département Sciences et Technologies, 97660 Dombeni, France*

² *UMR Espace-Dev, Université de Guyane, Campus de Troubiran 97300, Cayenne, France*

ARTICLE INFO

Article history:

Received: 10 June, 2017

Accepted: 14 July, 2017

Online: 28 December, 2017

Keywords:

Asymptotically periodic solution

Composition theorem

Evolutionary process

Stochastic evolution equation

ABSTRACT

In this paper, we study the existence (uniqueness) and asymptotic stability of the p -th mean S -asymptotically ω -periodic solutions for some non-autonomous Stochastic Evolution Equations driven by a \mathcal{Q} -Brownian motion. This is done using the Banach fixed point Theorem and a Gronwall inequality.

1 Introduction

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a complete probability space and $(\mathbb{H}, \|\cdot\|)$ a real separable Hilbert space. We are concerned in this paper with the existence and asymptotic stability of p -th mean S -asymptotically ω -periodic solution of the following stochastic evolution equation

$$\begin{cases} dX(t) = A(t)X(t)dt + f(t, X(t))dt \\ \quad + g(t, X(t))dW(t), & t \geq 0 \\ X(0) = c_0, \end{cases} \quad (1)$$

where $(A(t))_{t \geq 0}$ is a family of densely defined closed linear operators which generates an exponentially stable ω -periodic two-parameter evolutionary family. The functions $f : \mathbb{R}_+ \times L^p(\Omega, \mathbb{H}) \rightarrow \mathbb{L}^p(\Omega, \mathbb{H})$, $g : \mathbb{R}_+ \times \mathbb{L}^p(\Omega, \mathbb{H}) \rightarrow \mathbb{L}^p(\Omega, L_2^0)$ are continuous satisfying some additional conditions and $(W(t))_{t \geq 0}$ is a \mathcal{Q} -Brownian motion. The spaces $\mathbb{L}^p(\Omega, \mathbb{H})$, L_2^0 and the \mathcal{Q} -Brownian motion are defined in the next section.

The concept of periodicity is important in probability especially for investigations on stochastic processes. The interest in such a notion lies in its significance and applications arising in engineering, statistics, etc. In recent years, there has been an increasing interest in periodic solutions (pseudo-almost periodic, almost periodic, almost automorphic, asymptotically

almost periodic, etc) for stochastic evolution equations. For instance among others, let us mentioned the existence, uniqueness and asymptotic stability results of almost periodic solutions, almost automorphic solutions, pseudo almost periodic solutions studied by many authors, see, e.g. ([1]-[11]). The concept of S -asymptotically ω -periodic stochastic processes, which is the central question to be treated in this paper, was first introduced in the literature by Henriquez, Pierri et al in ([12, 13]). This notion has been developed by many authors.

In the literature, there has been a significant attention devoted this concept in the deterministic case; we refer the reader to ([14]-[20]) and the references therein. However, in the random case, there are few works related to the notion of S -asymptotically ω -periodicity with regard to the existence, uniqueness and asymptotic stability for stochastic processes. To our knowledge, the first work dedicated to S -asymptotically ω -periodicity for stochastic processes is due to S. Zhao and M. Song ([21, 22]) where they show existence of square-mean S -asymptotically ω -periodic solutions for a class of stochastic fractional functional differential equations and for a certain class of stochastic fractional evolution equation driven by Levy noise. But until now and to the best of our knowledge, there is no investigations for the exis-

*Corresponding Author: CUFR-Mayotte, IMAG-Montpellier, France, & solym.manou-abi@univ-mayotte.fr

tence (uniqueness), asymptotic stability of p -th mean S -asymptotically ω -periodic solutions when $p > 2$.

This paper is organized as follows. Section 2 deals with some preliminaries intended to clarify the presentation of concepts and norms used latter. We also give a composition result, see Theorem 1. In section 3 we present theoretical results on the existence and uniqueness of S -asymptotically ω -periodic solution of equation (1), see Theorem 2. We also present results on asymptotic stability of the unique S -asymptotically ω -periodic solution of equation(1), see Theorem 3.

2 Preliminaries

This section is concerned with some notations, definitions, lemmas and preliminary facts which are used in what follows.

2.1 p -th mean S asymptotically omega periodic process

Assume that the probability space $(\Omega, \mathcal{F}, \mathbb{P})$ is equipped with some filtration $(\mathcal{F}_t)_{t \geq 0}$ satisfying the usual conditions. Let $p \geq 2$. Denote by $\mathbb{L}^p(\Omega, \mathbb{H})$ the collection of all strongly measurable p -th integrable \mathbb{H} -valued random variables such that

$$\mathbb{E}\|X\|^p = \int_{\Omega} \|X(\omega)\|^p d\mathbb{P}(\omega) < \infty.$$

Definition 1 A stochastic process $X : \mathbb{R}_+ \rightarrow \mathbb{L}^p(\Omega, \mathbb{H})$ is said to be continuous whenever

$$\lim_{t \rightarrow s} \mathbb{E}\|X(t) - X(s)\|^p = 0.$$

Definition 2 A stochastic process $X : \mathbb{R}_+ \rightarrow \mathbb{L}^p(\Omega, \mathbb{H})$ is said to be bounded if there exists a constant $C > 0$ such that

$$\mathbb{E}\|X(t)\|^p \leq C \quad \forall t \geq 0$$

Definition 3 A continuous and bounded stochastic process $X : \mathbb{R}_+ \rightarrow \mathbb{L}^p(\Omega, \mathbb{H})$ is said to be p -mean S -asymptotically ω periodic if there exists $\omega > 0$ such that

$$\lim_{t \rightarrow +\infty} \mathbb{E}\|X(t + \omega) - X(t)\|^p = 0, \quad \forall t \geq 0.$$

The collection of p -mean S -asymptotically ω -periodic stochastic process with values in \mathbb{H} is then denoted by $SAP_{\omega}(\mathbb{L}^p(\Omega, \mathbb{H}))$.

A continuous bounded stochastic process X , which is 2-mean S -asymptotically ω -periodic is also called square-mean S -asymptotically ω -periodic.

Remark 1 Since any p -mean S -asymptotically ω -periodic process X is $\mathbb{L}^p(\Omega, \mathbb{H})$ bounded and continuous, the space $SAP_{\omega}(\mathbb{L}^p(\Omega, \mathbb{H}))$ is a Banach space equipped with the sup norm :

$$\|X\|_{\infty} = \sup_{t \geq 0} \left(\mathbb{E}\|X(t)\|^p \right)^{1/p}$$

Definition 4 A function $F : \mathbb{R}_+ \times \mathbb{L}^p(\Omega, \mathbb{H}) \rightarrow \mathbb{L}^p(\Omega, \mathbb{H})$ which is jointly continuous, is said to be p -mean S -asymptotically ω periodic in $t \in \mathbb{R}_+$ uniformly in $X \in K$ where $K \subseteq \mathbb{L}^p(\Omega, \mathbb{H})$ is bounded if for any $\epsilon > 0$ there exists $L_{\epsilon} > 0$ such that

$$\mathbb{E}\|F(t + \omega, X) - F(t, X)\|^p \leq \epsilon$$

for all $t \geq L_{\epsilon}$ and all process $X : \mathbb{R}_+ \rightarrow K$

Definition 5 A function $F : \mathbb{R}_+ \times \mathbb{L}^p(\Omega, \mathbb{H}) \rightarrow \mathbb{L}^p(\Omega, \mathbb{H})$ which is jointly continuous, is said to be p -mean asymptotically uniformly continuous on bounded sets $K' \subseteq \mathbb{L}^p(\Omega, \mathbb{H})$, if for all $\epsilon > 0$ there exists $\delta_{\epsilon} > 0$ such that

$$\mathbb{E}\|F(t, X) - F(t, Y)\|^p \leq \epsilon$$

for all $t \geq \delta_{\epsilon}$ and every $X, Y \in K'$ with $\mathbb{E}\|X - Y\|^p \leq \delta_{\epsilon}$.

Theorem 1 Let $F : \mathbb{R}_+ \times \mathbb{L}^p(\Omega, \mathbb{H}) \rightarrow \mathbb{L}^p(\Omega, \mathbb{H})$ be a p -mean S -asymptotically ω periodic in $t \in \mathbb{R}_+$ uniformly in $X \in K$ where $K \subseteq \mathbb{L}^p(\Omega, \mathbb{H})$ is bounded and p -mean asymptotically uniformly continuous on bounded sets. Assume that $X : \mathbb{R}_+ \rightarrow \mathbb{L}^p(\Omega, \mathbb{H})$ is a p -mean S asymptotically ω -periodic process. Then the stochastic process $(F(t, X(t)))_{t \geq 0}$ is p -mean S -asymptotically ω periodic.

Proof 1 Since $X : \mathbb{R}_+ \rightarrow \mathbb{L}^p(\Omega, \mathbb{H})$ is a p -mean S -asymptotically ω -periodic process, for all $\epsilon > 0$, there exists $T_{\epsilon} > 0$ such that for all $t \geq T_{\epsilon}$:

$$\mathbb{E}\|X(t + \omega) - X(t)\|^p \leq \epsilon \tag{2}$$

In addition X is bounded that is

$$\sup_{t \geq 0} \mathbb{E}\|X(t)\|^p < \infty$$

Let $K \subset \mathbb{L}^p(\Omega, \mathbb{H})$ be a bounded set such that $X(t) \in K$ for all $t \geq 0$.

We have :

$$\begin{aligned} \mathbb{E}\|F(t + \omega, X(t + \omega)) - F(t, X(t))\|^p & \\ & \leq 2^{p-1} \mathbb{E}\|F(t + \omega, X(t + \omega)) - F(t + \omega, X(t))\|^p \\ & \quad + 2^{p-1} \mathbb{E}\|F(t + \omega, X(t)) - F(t, X(t))\|^p \end{aligned}$$

Taking into account (2) and using the fact that F is p -mean asymptotically uniformly continuous on bounded sets, there exists $\delta_{\epsilon} = \epsilon$ and $L_{\epsilon} = T_{\epsilon}$ such that for all $t \geq T_{\epsilon}$:

$$\mathbb{E}\|F(t + \omega, X(t + \omega)) - F(t + \omega, X(t))\|^p \leq \frac{\epsilon}{2^p} \tag{3}$$

Similarly, using the p -mean S -asymptotically ω periodicity in $t \geq 0$ uniformly on bounded sets of F it follows that for all $t \geq T_{\epsilon}$:

$$\mathbb{E}\|F(t + \omega, X(t)) - F(t, X(t))\|^p \leq \frac{\epsilon}{2^p} \tag{4}$$

Bringing together the inequalities (3) and (4), we thus obtain that for all $t \geq T_{\epsilon} > 0$

$$\mathbb{E}\|F(t + \omega, X(t + \omega)) - F(t, X(t))\|^p \leq \epsilon$$

so that the stochastic process $t \rightarrow F(t, X(t))$ is p -mean S -asymptotically ω -periodic.

Lemma 1 Assume that $F : \mathbb{R}_+ \times \mathbb{L}^p(\Omega, \mathbb{H}) \rightarrow \mathbb{L}^p(\Omega, \mathbb{H})$ is p -mean uniformly S -asymptotically ω -periodic in $t \in \mathbb{R}_+$ uniformly on bounded sets and satisfies the Lipschitz condition, that is, there exists constant $L(F) > 0$ such that

$$\mathbb{E}\|F(t, X) - F(t, Y)\|^p \leq L(F)\mathbb{E}\|X - Y\|^p \quad \forall t \geq 0,$$

$\forall X, Y \in \mathbb{L}^p(\Omega, \mathbb{K})$. Let X be an p -mean S asymptotically ω -periodic proces, then the process $(F(t, X(t)))_{t \geq 0}$ is p -mean S -asymptotically ω -periodic.

For the proof, the reader can refer to [22] whenever $p = 2$. The case $p > 2$ is similar.

Now let us recall the notion of evolutionary family of operators.

Definition 6 A two-parameter family of bounded linear operators $\{U(t, s) : t \geq s \text{ with } t, s \geq 0\}$ from $\mathbb{L}^p(\Omega, \mathbb{H})$ into itself associate with $A(t)$ is called an evolutionary family of operators whenever the following conditions hold:

(a)

$$U(t, s)U(s, r) = U(t, r) \quad \text{for every } r \leq s \leq t;$$

(b)

$$U(t, t) = I, \text{ where } I \text{ is the identity operator;}$$

(c) For all $X \in \mathbb{L}^p(\Omega, \mathbb{H})$, the function $(t, s) \rightarrow U(t, s)X$ is continuous for $s < t$;

(d) The function $t \rightarrow U(t, s)$ is differentiable and

$$\frac{\partial}{\partial t}(U(t, s)) = A(t)U(t, s) \quad \text{for every } r \leq s \leq t;$$

For additional details on evolution families, we refer the reader to the book by Lunardi [23].

2.2 \mathcal{Q} -Brownian motion and Stochastic integrals

Let $(B_n(t))_{n \geq 1}, t \geq 0$ be a sequence of real valued standard Brownian motion mutully independent on the filtered space $(\Omega, \mathcal{F}, \mathbb{P}, \mathcal{F}_\square)$. Set

$$W(t) = \sum_{n \geq 1} \sqrt{\lambda_n} B_n(t) e_n, \quad t \geq 0,$$

where $\lambda_n \geq 0, n \geq 1$, are non negative real numbers and $(e_n)_{n \geq 1}$ the complete orthonormal basis in the Hilbert space $(\mathbb{H}, \|\cdot\|)$. Let \mathcal{Q} be a symmetric nonnegative operator with finite trace defined by

$$\mathcal{Q}e_n = \lambda_n e_n \quad \text{such that} \quad Tr(\mathcal{Q}) = \sum_{n \geq 1} \lambda_n < \infty$$

It is well known that $\mathbb{E}[W_t] = 0$ and for all $t \geq s \geq 0$, the distribution of $W(t) - W(s)$ is a Gaussian distribution $(\mathcal{N}(0, (t - s)\mathcal{Q}))$. The above-mentioned \mathbb{H} -valued stochastic process $(W(t))_{t \geq 0}$ is called an \mathcal{Q} -Brownian

motion.

Let $(\mathbb{K}, \|\cdot\|_{\mathbb{K}})$ be a real separable Hilbert space. Let also $\mathcal{L}(\mathbb{K}, \mathbb{H})$ be the space of all bounded linear operators from \mathbb{K} into \mathbb{H} . If $\mathbb{K} = \mathbb{H}$, we denote it by $\mathcal{L}(\mathbb{H})$.

Set $\mathbb{H}_0 = \mathcal{Q}^{1/2}\mathbb{H}$. The space \mathbb{H}_0 is a Hilbert space equipped with the norm $\|u\|_{\mathbb{H}_0} = \|\mathcal{Q}^{1/2}u\|$.

Define

$$L_2^0 = \{\Phi \in \mathcal{L}(\mathbb{H}_0, \mathbb{H}) : Tr[(\Phi\mathcal{Q}\Phi^*)] < \infty\}$$

the space of all Hilbert-Schmidt operators from \mathbb{H}_0 to \mathbb{H} equipped with the norm

$$\|\Phi\|_{L_2^0} = Tr[(\Phi\mathcal{Q}\Phi^*)] = \mathbb{E}\|\Phi\mathcal{Q}^{1/2}\|^2$$

In the sequel, to prove Lemma 4 and Theorem 2 we need the following Lemma that is a particular case of Lemma 2.2 in [24] (see also [25]).

Assume $T > 0$.

Lemma 2 Let $G : [0, T] \rightarrow \mathcal{L}(\mathbb{L}^p(\Omega, \mathbb{H}))$ be an \mathcal{F}_t -adapted measurable stochastic process satisfying $\int_0^T \mathbb{E}\|G(t)\|^2 dt < \infty$ almost surely. Then,

(i) the stochastic integral $\int_0^t G(s)dW(s)$ is a continuous, square integrable martingale with values in $(\mathbb{H}, \|\cdot\|)$ such that

$$\mathbb{E} \left\| \int_0^t G(s)dW(s) \right\|^2 \leq \mathbb{E} \int_0^t \|G(s)\|^2 ds$$

(ii) There exists some constant $C_p > 0$ such that the following particular case of Burkholder-Davis-Gundy inequality holds :

$$\mathbb{E} \sup_{0 \leq t \leq T} \left\| \int_0^t G(s)dW(s) \right\|^p \leq C_p \mathbb{E} \left(\int_0^T \|G(s)\|^2 ds \right)^{p/2}$$

In the sequel, we'll frequently make use of the following inequalities :

$$|a+b|^p \leq 2^{p-1}(|a|^p + |b|^p) \quad \text{for all } p \geq 1 \text{ and any real numbers } a, b.$$

$$\int_{t_0}^t e^{-2a(t-s)} ds \leq \int_{t_0}^t e^{-a(t-s)} ds \leq \frac{1}{a} \quad \forall t \geq t_0, \text{ where } a > 0.$$

3 Main results

In this section, we investigate the existence and the asymptotically stability of the p -th mean S -asymptotically ω -periodic solution to the already defined stochastic differential equation :

$$\begin{cases} dX(t) = A(t)X(t)dt + f(t, X(t))dt + g(t, X(t))dW(t), & t \geq 0 \\ X(0) = c_0, \end{cases} \quad (5)$$

where $A(t), t \geq 0$ is a family of densely defined closed linear operators and

$$f : \mathbb{R}_+ \times \mathbb{L}^p(\Omega, \mathbb{H}) \rightarrow \mathbb{L}^p(\Omega, \mathbb{H}),$$

$$g : \mathbb{R}_+ \times \mathbb{L}^p(\Omega, \mathbb{H}) \rightarrow \mathbb{L}^p(\Omega, L_2^0)$$

are jointly continuous satisfying some additional conditions and $(W(t))_{t \geq 0}$ is a \mathcal{Q} -Brownian motion with values in \mathbb{H} and \mathcal{F}_t -adapted.

Throughout the rest of this section, we require the following assumption on $U(t, s)$:

(H1): $A(t)$ generates an exponentially ω -periodic stable evolutionary process $(U(t, s))_{t \geq s}$ in $\mathbb{L}^p(\Omega, \mathbb{H})$, that is, a two-parameter family of bounded linear operators with the following additional conditions:

1. $U(t + \omega, s + \omega) = U(t, s)$ for all $t \geq s$ (ω -periodicity).
2. There exists $M > 0$ and $a > 0$ such that $\|U(t, s)\| \leq Me^{-a(t-s)}$ for $t \geq s$.

Now, note that if $A(t)$ generates an evolutionary family $(U(t, s))_{t \geq s}$ on $\mathbb{L}^p(\Omega, \mathbb{H})$ then the function g defined by $g(s) = U(t, s)X(s)$ where X is a solution of equation (1), satisfies the following relation

$$\begin{aligned} dg(s) &= -A(s)U(t, s)X(s) + U(t, s)dX(s) \\ &= -A(s)U(t, s)X(s) + A(s)U(t, s)X(s)ds \\ &\quad + U(t, s)f(s, X(s))ds + U(t, s)g(s, X(s))dW(s). \end{aligned}$$

Thus

$$dg(s) = U(t, s)f(s, X(s))ds + U(t, s)g(s, X(s))dW(s). \quad (6)$$

Integrating (6) on $[0, t]$ we obtain that

$$\begin{aligned} X(t) - U(t, 0)c_0 &= \int_0^t U(t, s)f(s, X(s))ds \\ &\quad + \int_0^t U(t, s)g(s, X(s))dW(s). \end{aligned}$$

Therefore, we define

Definition 7 An (\mathcal{F}_t) -adapted stochastic process $(X(t))_{t \geq 0}$ is called a mild solution of (1) if it satisfies the following stochastic integral equation:

$$\begin{aligned} X(t) &= U(t, 0)c_0 + \int_0^t U(t, s)f(s, X(s))ds \\ &\quad + \int_0^t U(t, s)g(s, X(s))dW(s). \end{aligned}$$

3.1 The existence of p -th mean S -asymptotically ω -periodic solution

We require the following additional assumptions:

(H.2) The function $f : \mathbb{R}_+ \times \mathbb{L}^p(\Omega, \mathbb{H}) \rightarrow \mathbb{L}^p(\Omega, \mathbb{H})$ is p -mean S -asymptotically ω periodic in $t \in \mathbb{R}_+$ uniformly in $X \in K$ where $K \subseteq \mathbb{L}^p(\Omega, \mathbb{H})$ is a bounded set. Moreover the function f satisfies the Lipschitz condition, that is, there exists constant $L(f) > 0$ such that

$$\mathbb{E}\|f(t, X) - f(t, Y)\|^p \leq L(f)\mathbb{E}\|X - Y\|^p \quad \forall t \geq 0$$

$\forall X, Y \in \mathbb{L}^p(\Omega, \mathbb{H})$.

(H.3) The function $g : \mathbb{R}_+ \times \mathbb{L}^p(\Omega, L_2^0) \rightarrow \mathbb{L}^p(\Omega, L_2^0)$ is p -mean S -asymptotically ω periodic in $t \in \mathbb{R}_+$ uniformly in $X \in K$ where $K \subseteq \mathbb{L}^p(\Omega, L_2^0)$ is a bounded set. Moreover the function g satisfies the Lipschitz condition, that is, there exists constant $L(g) > 0$ such that

$$\mathbb{E}\|g(t, X) - g(t, Y)\|_{L_2^0}^p \leq L(g)\mathbb{E}\|X - Y\|^p \quad \forall t \geq 0,$$

$\forall X, Y \in \mathbb{L}^p(\Omega, \mathbb{H})$.

Lemma 3 We assume that hypothesis **(H.1)** and **(H.2)** are satisfied. We define the nonlinear operator \wedge_1 by: for each $\phi \in SAP_\omega(\mathbb{L}^p(\Omega, \mathbb{H}))$

$$(\wedge_1 \phi)(t) = \int_0^t U(t, s)f(s, \phi(s))ds$$

Then the operator \wedge_1 maps $SAP_\omega(\mathbb{L}^p(\Omega, \mathbb{H}))$ into itself.

Proof 2 We define $h(s) = f(s, \phi(s))$. Since the hypothesis **(H.2)** is satisfied, using Lemma 1, we deduce that the function h is p -mean S -asymptotically ω -periodic.

Define $F(t) = \int_0^t U(t, s)h(s)ds$. It is easy to check that F is bounded and continuous. Now we have:

$$F(t + \omega) - F(t)$$

$$= \int_0^\omega U(t + \omega, s)h(s)ds + \int_0^t U(t, s)(h(s + \omega) - h(s))ds$$

$$= U(t + \omega, \omega) \int_0^\omega U(\omega, s)h(s)ds$$

$$+ \int_0^t U(t, s)(h(s + \omega) - h(s))ds$$

$$\mathbb{E}\|F(t + \omega) - F(t)\|^p$$

$$\leq 2^{p-1}M^{2p}e^{-apt}\mathbb{E}\left(\int_0^\omega e^{-a(\omega-s)}\|h(s)\|ds\right)^p$$

$$+ 2^{p-1}M^p\mathbb{E}\left(\int_0^t e^{-a(t-s)}\|h(s + \omega) - h(s)\|ds\right)^p$$

Let p and q be conjugate exponents. Using Hölder inequality, we obtain that $\mathbb{E}\|F(t + \omega) - F(t)\|^p$

$$\leq 2^{p-1}M^{2p}e^{-apt}\left(\int_0^\omega e^{-aq(\omega-s)}ds\right)^{p/q} \int_0^\omega \mathbb{E}\|h(s)\|^p ds$$

$$+ 2^{p-1}M^p\mathbb{E}\left(\int_0^t e^{-a(t-s)}\|h(s + \omega) - h(s)\|ds\right)^p$$

$$= I(t) + J(t)$$

where

$$I(t) = 2^{p-1}M^{2p}e^{-apt}\left(\int_0^\omega e^{-aq(\omega-s)}ds\right)^{p/q} \int_0^\omega \mathbb{E}\|h(s)\|^p ds$$

$$J(t) = 2^{p-1} M^p \mathbb{E} \left(\int_0^t e^{-a(t-s)} \|h(s+\omega) - h(s)\| ds \right)^p$$

$$= 2^{p-1} M^p \mathbb{E} \left(\int_0^t e^{-\frac{a}{q}(t-s)} \times e^{-\frac{a}{q}(t-s)} \|h(s+\omega) - h(s)\| ds \right)^p$$

It is obvious that

$$\lim_{t \rightarrow +\infty} J(t) = 0.$$

Using Hölder inequality, we obtain that $J(t)$

$$\leq 2^{p-1} M^p \left(\int_0^t e^{-a(t-s)} ds \right)^{p/q}$$

$$\int_0^t e^{-a(t-s)} \mathbb{E} \|h(s+\omega) - h(s)\|^p ds \quad (7)$$

$$\leq 2^{p-1} M^p \left(\frac{1}{a} \right)^{p/q} \int_0^t e^{-a(t-s)} \mathbb{E} \|h(s+\omega) - h(s)\|^p ds \quad (8)$$

$$(9)$$

Let $\epsilon > 0$. Since $\lim_{u \rightarrow +\infty} \mathbb{E} \|h(u+\omega) - h(u)\|^p = 0$:

$$\exists T_\epsilon > 0, u > T_\epsilon \Rightarrow \mathbb{E} \|h(u+\omega) - h(u)\|^p \leq \frac{\epsilon a^p}{2^{p-1} M^p}. \quad (10)$$

We have

$$J(t) \leq 2^{p-1} M^p \left(\frac{1}{a} \right)^{p/q} \int_0^{T_\epsilon} e^{-a(t-s)} \mathbb{E} \|h(s+\omega) - h(s)\|^p ds$$

$$+ 2^{p-1} M^p \left(\frac{1}{a} \right)^{p/q} \int_{T_\epsilon}^t e^{-a(t-s)} \mathbb{E} \|h(s+\omega) - h(s)\|^p ds$$

$$= J_1(t) + J_2(t),$$

where

$$J_1(t) = 2^{p-1} M^p \left(\frac{1}{a} \right)^{p/q} \int_0^{T_\epsilon} e^{-a(t-s)} \mathbb{E} \|h(s+\omega) - h(s)\|^p ds$$

$$J_2(t) = 2^{p-1} M^p \left(\frac{1}{a} \right)^{p/q} \int_{T_\epsilon}^t e^{-a(t-s)} \mathbb{E} \|h(s+\omega) - h(s)\|^p ds$$

Estimation of $J_1(t)$.

$$J_1(t) \leq 2^{p-1} M^p \left(\frac{1}{a} \right)^{p/q} \int_0^{T_\epsilon} e^{-a(t-s)} \mathbb{E} \|h(s+\omega) - h(s)\|^p ds$$

$$\leq 2^{p-1} M^p \left(\frac{1}{a} \right)^{p/q} 2^p \sup_{t \geq 0} \mathbb{E} \|h(t)\|^p e^{-at} \int_0^{T_\epsilon} e^{aqs} ds.$$

It is clear that $\lim_{t \rightarrow +\infty} J_1(t) = 0$.

Estimation of $J_2(t)$.

Using the Inequality in (10) we have

$$J_2(t) = 2^{p-1} M^p \left(\frac{1}{a} \right)^{p/q} \int_{T_\epsilon}^t e^{-a(t-s)} \mathbb{E} \|h(s+\omega) - h(s)\|^p ds$$

$$\leq 2^{p-1} M^p \left(\frac{1}{a} \right)^{p/q} \left(\frac{1}{a} \right) \frac{\epsilon a^p}{2^{p-1} M^p}$$

$$= 2^{p-1} M^p a^{-p} \frac{\epsilon a^p}{2^{p-1} M^p}$$

$$\leq \epsilon.$$

Lemma 4 We assume that hypothesis **(H.1)** and **(H.3)** are satisfied. We define the nonlinear operator Λ_2 by: for each $\phi \in SAP_\omega(\mathbb{L}^p(\Omega, L_2^0))$

$$(\Lambda_2 \phi)(t) = \int_0^t U(t,s) g(s, \phi(s)) dW(s)$$

Then the operator Λ_2 maps $SAP_\omega(\mathbb{L}^p(\Omega, L_2^0))$ into itself.

Proof 3 We define $h(s) = g(s, \phi(s))$. Since the hypothesis **(H.3)** is satisfied, using Lemma 1, we deduce that the function h is p -mean S asymptotically ω periodic.

Define $F(t) = \int_0^t U(t,s) h(s) dW(s)$. It is easy to check that F is bounded and continuous. We have :

$$F(t+\omega) - F(t)$$

$$= \int_0^\omega U(t+\omega, s) h(s) dW(s) + \int_0^t U(t, s) (h(s+\omega) - h(s)) dW(s)$$

$$= U(t+\omega, \omega) \int_0^\omega U(\omega, s) h(s) dW(s) + \int_0^t U(t, s) (h(s+\omega) - h(s)) dW(s)$$

$$= U(t+\omega, \omega) F(\omega) + \int_0^t U(t, s) (h(s+\omega) - h(s)) ds$$

$$\mathbb{E} \|F(t+\omega) - F(t)\|^p$$

$$\leq 2^{p-1} M^p e^{-apt} \mathbb{E} \left\| \int_0^\omega U(\omega, s) h(s) dW(s) \right\|^p$$

$$+ 2^{p-1} \mathbb{E} \left\| \int_0^t U(t, s) (h(s+\omega) - h(s)) dW(s) \right\|^p$$

$$:= I(t) + J(t)$$

where

$$I(t) = 2^{p-1} M^p e^{-pat} \mathbb{E} \left\| \int_0^\omega U(\omega, s) h(s) dW(s) \right\|^p$$

$$J(t) = 2^{p-1} \mathbb{E} \left\| \int_0^t U(t, s) (h(s+\omega) - h(s)) dW(s) \right\|^p$$

It is clear that

$$\lim_{t \rightarrow +\infty} \mathbb{E} I(t) = 0.$$

Let $\epsilon > 0$. Since $\lim_{t \rightarrow +\infty} \mathbb{E} \|h(t+\omega) - h(t)\|^p = 0$

$$\exists T_\epsilon > 0, t > T_\epsilon \Rightarrow \mathbb{E} \|h(s+\omega) - h(s)\|_{L_2^0}^p \leq \frac{\epsilon (2a)^{p/2}}{4^{p-1} M^p C_p^p}, \quad (11)$$

where the constant C_p will be precised in the next lines. We have $\mathbb{E}J(t)$

$$\begin{aligned} &= 2^{p-1} \mathbb{E} \left\| \int_0^t U(t,s)(h(s+\omega) - h(s)) dW(s) \right\|^p \\ &\leq 4^{p-1} \mathbb{E} \left\| \int_0^{T_\epsilon} U(t,s)(h(s+\omega) - h(s)) dW(s) \right\|^p \\ &+ 4^{p-1} \mathbb{E} \left\| \int_{T_\epsilon}^t U(t,s)(h(s+\omega) - h(s)) dW(s) \right\|^p \\ &:= \mathbb{E}J_1(t) + \mathbb{E}J_2(t), \end{aligned}$$

where

$$\begin{aligned} J_1(t) &= 4^{p-1} \left\| \int_0^{T_\epsilon} U(t,s)(h(s+\omega) - h(s)) dW(s) \right\|^p \\ J_2(t) &= 4^{p-1} \left\| \int_{T_\epsilon}^t U(t,s)(h(s+\omega) - h(s)) dW(s) \right\|^p \end{aligned}$$

Note that for all $t \geq 0$,

$$h(t+\omega) - h(t) \in \mathbb{L}^p(\Omega, L_2^0) \subseteq \mathbb{L}^2(\Omega, L_2^0)$$

and

$$\begin{aligned} &\int_0^t \mathbb{E} \|U(t,s)(h(s+\omega) - h(s))\|^2 ds \\ &\leq M^2 \int_0^t e^{-2a(t-s)} \mathbb{E} \|h(s+\omega) - h(s)\|_{L_2^0}^2 ds \\ &\leq 4M^2 \sup_{t \geq 0} \mathbb{E} \|h(t)\|_{L_2^0}^2 \int_0^t e^{-2a(t-s)} ds \\ &\leq 4M^2 a^{-2} \sup_{t \geq 0} \mathbb{E} \|h(t)\|_{L_2^0}^2 \\ &< \infty. \end{aligned}$$

Estimation of $\mathbb{E}J_1(t)$.

Assume that $p > 2$. Using Hölder inequality between conjugate exponents $\frac{p}{p-2}$ and $\frac{p}{2}$ together with Lemma 2, part (ii), there exists constant C_p such that : $\mathbb{E}J_1(t)$

$$\begin{aligned} &\leq C_p 4^{p-1} M^p \mathbb{E} \left[\int_0^{T_\epsilon} e^{-2a(t-s)} \|h(s+\omega) - h(s)\|_{L_2^0}^2 ds \right]^{p/2} \\ &\leq C_p 4^{p-1} M^p \left(\int_0^{T_\epsilon} e^{-\frac{2ap(t-s)}{p-2}} ds \right)^{\frac{p-2}{2}} \\ &\quad \int_0^{T_\epsilon} \mathbb{E} \|h(s+\omega) - h(s)\|_{L_2^0}^p ds \\ &\leq C_p 4^{p-1} M^p e^{-apt} \left(\int_0^{T_\epsilon} e^{\frac{2aps}{p-2}} ds \right)^{\frac{p-2}{2}} T_\epsilon 2^p \sup_{s \geq 0} \mathbb{E} \|h(s)\|_{L_2^0}^p \end{aligned}$$

Therefore

$$\lim_{t \rightarrow +\infty} \mathbb{E}J_1(t) = 0.$$

Assume that $p = 2$. By Lemma 2, part (i) we get : $\mathbb{E}J_1(t)$

$$\begin{aligned} &\leq 4M^2 \mathbb{E} \left[\int_0^{T_\epsilon} e^{-a(t-s)} \|h(s+\omega) - h(s)\|_{L_2^0} dB(s) \right]^2 \\ &\leq 4M^2 \mathbb{E} \left[\int_0^{T_\epsilon} e^{-2a(t-s)} \|h(s+\omega) - h(s)\|_{L_2^0}^2 ds \right] \\ &\leq 16M^2 e^{-2at} \sup_{s \geq 0} \mathbb{E} \|h(s)\|_{L_2^0}^2 \int_0^{T_\epsilon} e^{2as} ds \\ &\leq 16M^2 e^{-2at} \sup_{s \geq 0} \mathbb{E} \|h(s)\|_{L_2^0}^2 \int_0^{T_\epsilon} e^{2as} ds \end{aligned}$$

Thus

$$\lim_{t \rightarrow +\infty} \mathbb{E}J_1(t) = 0.$$

Estimation of $\mathbb{E}J_2(t)$.

Assume that $p > 2$. Using again Lemma 2, part (ii), Hölder inequality between conjugate exponents $\frac{p}{p-2}$ and $\frac{p}{2}$ and the inequality in (11) we have $\mathbb{E}J_2(t)$

$$\begin{aligned} &= 4^{p-1} \mathbb{E} \left\| \int_{T_\epsilon}^t U(t,s)(h(s+\omega) - h(s)) dW(s) \right\|^p \\ &\leq 4^{p-1} M^p C_p \mathbb{E} \left[\int_{T_\epsilon}^t e^{-2a(t-s)} \|h(s+\omega) - h(s)\|_{L_2^0}^2 ds \right]^{p/2} \\ &= 4^{p-1} M^p C_p \mathbb{E} \left[\int_{T_\epsilon}^t e^{-2a(t-s)\frac{p-2}{p}} \times e^{-2a(t-s)\frac{2}{p}} \|h(s+\omega) - h(s)\|_{L_2^0}^2 ds \right]^{p/2} \\ &\leq C_p 4^{p-1} M^p \left(\int_{T_\epsilon}^t e^{-2a(t-s)} ds \right)^{\frac{p-2}{2}} \int_{T_\epsilon}^t e^{-2a(t-s)} \mathbb{E} \|h(s+\omega) - h(s)\|_{L_2^0}^p ds \\ &\leq \frac{C_p 4^{p-1} M^p \epsilon (2a)^{p/2}}{C_p 4^{p-1} M^p} \left(\int_{T_\epsilon}^t e^{-2a(t-s)} ds \right)^{\frac{p}{2}} \\ &\leq \epsilon. \end{aligned}$$

We conclude that

$$\lim_{t \rightarrow +\infty} \mathbb{E}J_2(t) = 0.$$

Assume that $p = 2$. By Lemma 2, part (i) and Cauchy-Schwarz inequality we have $\mathbb{E}J_2(t)$

$$\begin{aligned} &= 4 \mathbb{E} \left\| \int_{T_\epsilon}^t U(t,s)(h(s+\omega) - h(s)) dW(s) \right\|^2 \\ &\leq 4M^2 \mathbb{E} \left[\int_{T_\epsilon}^t e^{-2a(t-s)} \|h(s+\omega) - h(s)\|_{L_2^0}^2 ds \right] \\ &= 4M^2 \left[\int_{T_\epsilon}^t e^{-a(t-s)} \times e^{-a(t-s)} \mathbb{E} \|h(s+\omega) - h(s)\|_{L_2^0}^2 ds \right] \\ &\leq 4M^2 \left(\int_{T_\epsilon}^t e^{-2a(t-s)} ds \right)^{1/2} \\ &\quad \left(\int_{T_\epsilon}^t e^{-2a(t-s)} (\mathbb{E} \|h(s+\omega) - h(s)\|_{L_2^0}^2)^2 ds \right)^{1/2} \end{aligned}$$

Note also that for $t \geq T_\epsilon$:

$$\mathbb{E} \|h(s+\omega) - h(s)\|_{L_2^0}^2 \leq \frac{\epsilon a}{2M^2}$$

so that

$$\mathbb{E}J_2(t) \leq \frac{4M^2\epsilon a}{2M^2} \left(\int_{T_\epsilon}^t e^{-2a(t-s)} ds \right) \leq \epsilon.$$

This implies that

$$\lim_{t \rightarrow +\infty} \mathbb{E}J_2(t) = 0$$

Finally, we conclude that

$$\lim_{t \rightarrow +\infty} \mathbb{E}\|F(t + \omega) - F(t)\|^p = 0$$

Theorem 2 We assume that hypothesis (H.1), (H.2) and (H.3) are satisfied and

(i)

$$\Theta = 2^{p-1}M^p \left(L(f)a^{-p} + C_p L(g)a^{-\frac{p}{2}} \right) < 1 \quad \text{if } p > 2 \quad (12)$$

(ii)

$$\Xi = 2M^2 \left(L(f)\frac{1}{a^2} + L(g)\frac{1}{a} \right) < 1 \quad \text{if } p = 2 \quad (13)$$

Then the stochastic evolution equation (1) has a unique p -mean S -asymptotically ω -periodic solution.

Proof 4 We define the nonlinear operator Γ by the expression

$$(\Gamma\Phi)(t) = U(t, 0)c_0 +$$

$$\int_0^t U(t, s)f(s, \Phi(s))ds + \int_0^t U(t, s)g(s, \Phi(s))dW(s)$$

Note that

$$(\Gamma\Phi)(t) = U(t, 0)c_0 + (\Lambda_1\Phi)(t) + (\Lambda_2\Phi)(t)$$

According to the hypothesis (H1) we have :

$$\mathbb{E}\|U(t + \omega, 0) - U(t, 0)\|^p$$

$$\leq 2^{p-1} \left(\mathbb{E}\|U(t + \omega, 0)\|^p + \mathbb{E}\|U(t, 0)\|^p \right)$$

$$\leq 2^{p-1}M^p e^{-ap(t+\omega)} + M^p e^{-apt}$$

$$= 2^{p-1}M^p e^{-apt} (e^{-ap\omega} + 1)$$

Therefore

$$\lim_{t \rightarrow +\infty} \mathbb{E}\|U(t + \omega, 0) - U(t, 0)\|^p = 0$$

According to Lemma 3 and Lemma 4, the operators Λ_1 and Λ_2 maps the space of p -mean S -asymptotically ω -periodic solutions into itself. Thus Γ maps the space of p -mean S -asymptotically ω periodic solutions into itself.

We have

$$\mathbb{E}\|\Gamma\Phi(t) - \Gamma\Psi(t)\|^p$$

$$\leq 2^{p-1} \mathbb{E} \left(\int_0^t \|U(t, s)\| \|f(s, \Phi(s)) - f(s, \Psi(s))\| ds \right)^p$$

$$+ 2^{p-1} \mathbb{E} \left(\int_0^t \|U(t, s)\| \|g(s, \Phi(s)) - g(s, \Psi(s))\| dW(s) \right)^p$$

$$\leq 2^{p-1}M^p \mathbb{E} \left(\int_0^t e^{-a(t-s)} \|f(s, \Phi(s)) - f(s, \Psi(s))\| ds \right)^p$$

$$+ 2^{p-1}M^p \mathbb{E} \left(\int_0^t e^{-a(t-s)} \|g(s, \Phi(s)) - g(s, \Psi(s))\| dW(s) \right)^p$$

Case $p > 2$: By Lemma 2, part (ii) and Hölder inequality we have

$$\mathbb{E}\|\Gamma\Phi(t) - \Gamma\Psi(t)\|^p$$

$$\leq 2^{p-1}M^p L(f) \left(\int_0^t e^{-a(t-s)} ds \right)^{p-1} \int_0^t e^{-a(t-s)} \mathbb{E}\|\Phi(s) - \Psi(s)\|^p ds$$

$$+ 2^{p-1}M^p C_p \mathbb{E} \left(\int_0^t e^{-2a(t-s)} \|g(s, \Phi(s)) - g(s, \Psi(s))\|_{L_2^0}^2 ds \right)^{p/2}$$

$$\leq 2^{p-1}M^p L(f) \sup_{s \geq 0} \mathbb{E}\|\Phi(s) - \Psi(s)\|^p \left(\int_0^t e^{-a(t-s)} ds \right)^p$$

$$+ 2^{p-1}M^p C_p \mathbb{E} \left(\int_0^t e^{-a(t-s)} \|g(s, \Phi(s)) - g(s, \Psi(s))\|_{L_2^0}^2 ds \right)^{p/2}$$

$$\leq 2^{p-1}M^p a^{-p} L(f) \|\Phi - \Psi\|_\infty^p +$$

$$2^{p-1}M^p C_p \mathbb{E} \left(\int_0^t e^{-\frac{ap(t-s)}{p-2}} e^{-\frac{a(t-s)}{p}} \|g(s, \Phi(s)) - g(s, \Psi(s))\|_{L_2^0}^2 ds \right)^{p/2}$$

$$\leq 2^{p-1}M^p a^{-p} L(f) \|\Phi - \Psi\|_\infty^p$$

$$+ 2^{p-1}M^p C_p \left(\int_0^t e^{-a(t-s)} ds \right)^{\frac{p-2}{2}} \times$$

$$\int_0^t e^{-a(t-s)} \mathbb{E}\|g(s, \Phi(s)) - g(s, \Psi(s))\|_{L_2^0}^p ds$$

$$\leq 2^{p-1}M^p \left(L(f)a^{-p} + C_p L(g)a^{-\frac{p}{2}} \right) \|\Phi - \Psi\|_\infty^p$$

This implies that

$$\|\Gamma\Phi - \Gamma\Psi\|_\infty^p \leq 2^{p-1}M^p \left(L(f)a^{-p} + C_p L(g)a^{-\frac{p}{2}} \right) \|\Phi - \Psi\|_\infty^p$$

Consequently, if $\Theta < 1$, then Γ is a contraction mapping.

One completes the proof by the Banach fixed-point principle.

Case $p = 2$: using Cauchy-Schwarz inequality and Lemma 2, part (i), we obtain

$$\mathbb{E}\|\Gamma\Phi(t) - \Gamma\Psi(t)\|^2$$

$$\leq 2M^2 \left(\int_0^t e^{-a(t-s)} ds \right) \mathbb{E} \int_0^t e^{-a(t-s)} \|f(s, \Phi(s)) - f(s, \Psi(s))\|^2 ds$$

$$+ 2M^2 \mathbb{E} \int_0^t e^{-2a(t-s)} \|g(s, \Phi(s)) - g(s, \Psi(s))\|_{L_2^0}^2 ds$$

$$\leq 2M^2 L(f) \sup_{s \geq 0} \mathbb{E}\|\Phi(s) - \Psi(s)\|^2 \left(\int_0^t e^{-a(t-s)} ds \right)^2$$

$$+ 2M^2 L(g) \sup_{s \geq 0} \mathbb{E}\|\Phi(s) - \Psi(s)\|^2 \int_0^t e^{-2a(t-s)} ds$$

$$\leq 2M^2 \left(L(f)\frac{1}{a^2} + L(g)\frac{1}{a} \right) \sup_{s \geq 0} \mathbb{E}\|\Phi(s) - \Psi(s)\|^2$$

This implies that

$$\|\Gamma\Phi - \Gamma\Psi\|_\infty^2 \leq 2M^2 \left(L(f) \frac{1}{a^2} + L(g) \frac{1}{a} \right) \|\Phi - \Psi\|_\infty^2$$

Consequently, if $\Xi < 1$, then Γ is a contraction mapping. One completes the proof by the Banach fixed-point principle.

3.2 Stability of p -mean S asymptotically ω periodic solution

In the previous section, for the non linear SDE, we obtain that it has a unique p -mean S-asymptotically ω -periodic solution under some conditions. In this section, we will show that the unique p mean S asymptotically ω periodic solution is asymptotically stable in the p mean sense.

Recall that

Definition 8 The unique p -mean S asymptotically ω periodic solution $X^*(t)$ of (1) is said to be stable in p -mean sense if for any $\epsilon > 0$, there exists $\delta > 0$ such that

$$\mathbb{E}\|X(t) - X^*(t)\|^p < \epsilon, \quad t \geq 0,$$

whenever $\mathbb{E}\|X(0) - X^*(0)\|^p < \delta$, where $X(t)$ stands for a solution of (1) with initial value $X(0)$.

Definition 9 The unique p -mean S asymptotically ω periodic solution $X^*(t)$ is said to be asymptotically stable in p -mean sense if it is stable in p -mean sense and

$$\lim_{t \rightarrow \infty} \mathbb{E}\|X(t) - X^*(t)\|^p = 0$$

The following Gronwall inequality is proved to be useful in our asymptotical stability analysis.

Lemma 5 Let $u(t)$ be a non negative continuous functions for $t \geq 0$, and α, γ be some positive constants. If

$$u(t) \leq \alpha e^{-\beta t} + \gamma \int_0^t e^{-\beta(t-s)} u(s) ds, \quad t \geq 0,$$

then

$$u(t) \leq \alpha \exp\{(-\beta + \gamma)t\}$$

Theorem 3 Suppose that hypothesis (H.1), (H.2) and (H.3) are satisfied and assume that

$$(i) \quad 3^{p-1} M^p \left(L(f) a^{1-p} + L(g) C_p a^{\frac{2-p}{2}} \right) < a \quad (14)$$

whenever $p > 2$.

$$(ii) \quad 3M^2 \left(L(f) a^{-1} + L(g) \right) < a \quad (15)$$

whenever $p = 2$.

Then the p -mean S-asymptotically solution X_t^* of (1) is asymptotically stable in the p -mean sense.

Remark 2 Note that the above conditions (14) respectively (15) implies conditions (12) respectively (13) of Theorem 2.

Proof 5 $\mathbb{E}\|X(t) - X^*(t)\|^p$

$$\begin{aligned} &= \mathbb{E} \left\| U(t, 0)(X(0) - X^*(0)) \right. \\ &\quad + \int_0^t U(t, s) (f(s, X(s)) - f(s, X^*(s))) ds \\ &\quad \left. + \int_0^t U(t, s) (g(s, X(s)) - g(s, X^*(s))) dW(s) \right\|^p \end{aligned}$$

Assume that $p > 2$. Using Hölder inequality we have $\mathbb{E}\|X(t) - X^*(t)\|^p$

$$\begin{aligned} &\leq 3^{p-1} M^p e^{-ap t} \mathbb{E}\|X(0) - X^*(0)\|^p \\ &\quad + 3^{p-1} \mathbb{E} \left(\int_0^t \|U(t, s)\| \|f(s, X(s)) - f(s, X^*(s))\| ds \right)^p \\ &\quad + 3^{p-1} \mathbb{E} \left(\int_0^t \|U(t, s)\| \|g(s, X(s)) - g(s, X^*(s))\| dW(s) \right)^p \\ &\leq 3^{p-1} M^p e^{-ap t} \mathbb{E}\|X(0) - X^*(0)\|^p \\ &\quad + 3^{p-1} M^p \mathbb{E} \left(\int_0^t e^{-a(t-s)} \|f(s, X(s)) - f(s, X^*(s))\| ds \right)^p \\ &\quad + 3^{p-1} M^p \mathbb{E} \left(\int_0^t e^{-a(t-s)} \|g(s, X(s)) - g(s, X^*(s))\| dW(s) \right)^p \\ &= 3^{p-1} M^p e^{-ap t} \mathbb{E}\|X(0) - X^*(0)\|^p + 3^{p-1} M^p \times \left(\mathbb{E} \left(\int_0^t e^{-\frac{a(p-1)(t-s)}{p}} e^{-\frac{a(t-s)}{p}} \|f(s, X(s)) - f(s, X^*(s))\| ds \right)^p \right) \\ &\quad + 3^{p-1} M^p C_p \times \left(\int_0^t e^{-\frac{2a(p-2)(t-s)}{p}} e^{-\frac{4a(t-s)}{p}} \mathbb{E}\|g(s, X(s)) - g(s, X^*(s))\|_{L_2^0}^2 ds \right)^{p/2} \\ &\leq 3^{p-1} M^p e^{-ap t} \mathbb{E}\|X(0) - X^*(0)\|^p \\ &\quad + 3^{p-1} M^p L(f) \left(\int_0^t e^{-a(t-s)} ds \right)^{p-1} \times \left(\int_0^t e^{-a(t-s)} \mathbb{E}\|X(s) - X^*(s)\|^p ds \right) \\ &\quad + 3^{p-1} M^p C_p \left(\int_0^t e^{-2a(t-s)} \right)^{\frac{p-2}{2}} \times \left(\int_0^t e^{-2a(t-s)} \mathbb{E}\|g(s, X(s)) - g(s, X^*(s))\|_{L_2^0}^p ds \right) \end{aligned}$$

so that

$$\begin{aligned} &\mathbb{E}\|X(t) - X^*(t)\|^p \\ &\leq 3^{p-1} M^p e^{-ap t} \mathbb{E}\|X(0) - X^*(0)\|^p \\ &\quad + 3^{p-1} M^p L(f) \left(\frac{1}{a} \right)^{p-1} \int_0^t e^{-a(t-s)} \mathbb{E}\|X(s) - X^*(s)\|^p ds \\ &\quad + 3^{p-1} M^p C_p L(g) \left(\frac{1}{a} \right)^{\frac{p-2}{2}} \int_0^t e^{-a(t-s)} \mathbb{E}\|X(s) - X^*(s)\|^p ds \end{aligned}$$

Using Lemma 5, we obtain :

$$\begin{aligned} &\mathbb{E}\|X(t) - X^*(t)\|^p \\ &\leq 3^{p-1} M^p \times \mathbb{E}\|X(0) - X^*(0)\|^p \times \\ &\quad \exp \left\{ \left(-a + 3^{p-1} M^p \left(L(f) a^{1-p} + L(g) C_p a^{\frac{2-p}{2}} \right) \right) t \right\} \end{aligned}$$

Straightforwardly, we obtain that $X(t)$ converges to 0 exponentially fast if

$$-a + 3^{p-1} M^p \left(L(f) \left(\frac{1}{a} \right)^{p-1} + L(g) C_p a^{\frac{2-p}{2}} \right) < 0,$$

which is equivalent to our condition (14). Therefore X^* is asymptotically stable in the p -mean sense.

Assume that $p = 2$. We have

$$\begin{aligned} & \mathbb{E} \|X(t) - X^*(t)\|^2 \\ & \leq 3M^2 e^{-at} \mathbb{E} \|X(0) - X^*(0)\|^2 \\ & \quad + 3\mathbb{E} \left(\int_0^t \|U(t,s)\| \|f(s, X(s)) - f(s, X^*(s))\| ds \right)^2 \\ & \quad + 3\mathbb{E} \left(\int_0^t \|U(t,s)\| \|g(s, X(s)) - g(s, X^*(s))\| dW(s) \right)^2 \\ & \leq 3M^2 e^{-at} \mathbb{E} \|X(0) - X^*(0)\|^2 \\ & \quad + 3M^2 \mathbb{E} \left(\int_0^t e^{-a(t-s)} \|f(s, X(s)) - f(s, X^*(s))\| ds \right)^2 \\ & \quad + 3M^2 \mathbb{E} \left(\int_0^t e^{-a(t-s)} \|g(s, X(s)) - g(s, X^*(s))\| dW(s) \right)^2 \end{aligned}$$

Then using Cauchy-Schwartz inequality and Lemma 2.3 part (i), we have

$$\begin{aligned} & \mathbb{E} \|X(t) - X^*(t)\|^2 \\ & \leq 3M^2 e^{-at} \mathbb{E} \|X(0) - X^*(0)\|^2 \\ & \quad + 3M^2 \int_0^t e^{-a(t-s)} ds \int_0^t e^{-a(t-s)} \mathbb{E} \|f(s, X(s)) - f(s, X^*(s))\|^2 ds \\ & \quad + 3M^2 \int_0^t e^{-2a(t-s)} \mathbb{E} \|g(s, X(s)) - g(s, X^*(s))\|_{L_2^0}^2 ds, \\ & \leq 3M^2 e^{-at} \mathbb{E} \|X(0) - X^*(0)\|^2 \\ & \quad + 3M^2 L(f) a^{-1} \int_0^t e^{-a(t-s)} \mathbb{E} \|X(s) - X^*(s)\|^2 ds \\ & \quad + 3M^2 L(g) \int_0^t e^{-a(t-s)} \mathbb{E} \|X(s) - X^*(s)\|^2 ds \end{aligned}$$

Thus

$$\begin{aligned} & \mathbb{E} \|X(t) - X^*(t)\|^2 \\ & \leq 3M^2 e^{-at} \mathbb{E} \|X(0) - X^*(0)\|^2 \\ & \quad + \left(\frac{3M^2 L(f)}{a} + 3M^2 L(g) \right) \int_0^t e^{-a(t-s)} \mathbb{E} \|X(s) - X^*(s)\|^2 ds \end{aligned}$$

By Lemma 5 we have

$$\begin{aligned} & \mathbb{E} \|X(t) - X^*(t)\|^2 \\ & \leq 3M^2 \mathbb{E} \|X(0) - X^*(0)\|^2 \exp \left\{ \left(-a + 3M^2 \left(\frac{L(f)}{a} + L(g) \right) \right) t \right\} \end{aligned}$$

Therefore $\mathbb{E} \|X(t) - X^*(t)\|^2$ converges to 0 exponentially fast whenever condition (15) holds. In particular the unique S -asymptotically ω -periodic solution is asymptotically stable in square mean sense.

Acknowledgement

The first author thanks the Centre Universitaire de Mayotte for the financial support. The authors declares that there is no conflict of interest regarding the publication of this paper.

References

- [1] P. Bezandry and T. Diagana, *Existence of Almost Periodic Solutions to Some Stochastic Differential Equations*. *Applicable Analysis*, 86 (7), 819-827.
- [2] P. Bezandry and T. Diagana, *Square-Mean Almost Periodic Solutions Nonautonomous Stochastic Differential Equations*. *Electron. J. Differential Equations* (2007), no. 117, 1-10.
- [3] P. Bezandry and T. Diagana, *Existence of Quadratic-Mean Almost Periodic Solutions to Some Stochastic Hyperbolic Differential Equations*. *Electron. J. Differential Equations*, 2009, vol. 2009, no. 111, 1-14.
- [4] P. Bezandry and T. Diagana, *Square-Mean Almost Periodic solutions to Some Stochastic Hyperbolic Differential Equations with Infinite Delay*. *Commun. Math. Anal.* 8 (2010), no.2, 1-22
- [5] P. Bezandry and T. Diagana, *p-th Mean Pseudo Almost Automorphic Mild solutions to Some Nonautonomous Stochastic Differential Equations*. *Afr. Diaspora. J. Math.* Vol. 12 (2011), no.1, 60-79
- [6] J. Cao, Q. Yang, Z. Huang and Q. Liu *Asymptotically almost periodic solutions of stochastic functional differential equations*. *Applied Mathematics and Computation* 218 (2011) 1499-1511
- [7] Y-K. Chang, Z-H. Zhao and G.M. N'Gurkaka *Square mean almost automorphic mild solutions to non-autonomous stochastic differential equations in Hilbert spaces*. *Computers and Mathematics with Applications* 61 (2011), 384-391.
- [8] M.A. Diop, K.Ezzinbi and M.M. Mbaye. *Existence and global attractiveness of a pseudo almost periodic solution in p-th mean sense for stochastic evolution equation driven by a fractional Brownian motion*. *Stochastics An International Journal of Probability and Stochastic Processes* 87(6):1-33, 2015.
- [9] Xie Liang, *Square-Mean Almost Periodic solutions to Some Stochastic Evolutions Equations*. *Acta Mathematica Sinica. English Series.* (2014) Vol. 30, no.5, 881-898
- [10] ZhenXin Liu and Kai Sun. *Almost automorphic solutions to SDE driven by Levy noise*. *Journal of Functional Analysis* 266 (2014), 1115-1149
- [11] M. Zhang and G. Zong. *Almost Periodic Solutions for stochastic Differential Equations driven by G-Brownian motion*. *Communications in Statistics. Theory and Methods.* 44(11), 2371-2384 (2015)
- [12] H. R. Henríquez, M. Pierri and P. Táboas. *On S-asymptotically ω -periodic function on Banach spaces and applications*. *J. Math. Anal Appl.* 343, 1119-1130, 2008.
- [13] H. R. Henríquez, M. Pierri and P. Táboas. *Existence of S-asymptotically ω -periodic solutions for abstract neutral equations*. *Bull. Aust. Math. Soc.* 78, 365-382, 2008.
- [14] J. Blot, P. Cieutat and G. M. N'Guérékata *S-asymptotically ω -periodic functions and applications to evolution equations*. *African Diaspora J. Math.* 12, 113-121, 2011.
- [15] C. Cuevas and J.C. de Souza. *S-asymptotically ω -periodic solutions of semilinear fractional integro-differential equations*. *Appl. Math. Lett.* 22, 865-870, 2009.
- [16] C. Cuevas and C. Lizanna. *S-asymptotically ω -periodic solutions of semilinear Volterra equations*. *Math. Meth. Appl. Sci.* 33, 1628-1636, 2010.

- [17] C. Cuevas and C. Lizanna. *Existence of S-asymptotically ω -periodic solutions for two-times fractional order differential equations*. *Southeast.Asian Bull.Math.* 37, 683-690, 2013.
- [18] G. Da Prato and J. Zabczyk. *Stochastic Equations in Infinite Dimensions*. *Encyclopedia of Mathematics and Its Applications*, 44, Cambridge University Press, Cambridge, 1992.
- [19] W. Dimbour J-C. Mado. *S-asymptotically ω -periodic solution for a nonlinear differential equation with piecewise constant argument in a Banach space*. *CUBO A Mathematical Journal* 16(13), 55-65, 2014.
- [20] W. Dimbour, G.Mophou and G.M. N'Guérékata. *S-asymptotically ω -periodic solution for partial differential equations with finite delay*. *Electron.J.Differ.Equa.* 2011, 1-12, 2011.
- [21] Shufen Zhao and Minghui Song. *S-asymptotically ω -periodic solutions in distribution for a class of Stochastic fractional functional differential equations*. *arXiv : 1609.01453v1 [math.DS]*. 6 Sep 2016.
- [22] Shufen Zhao and Minghui Song. *Square-mean S-asymptotically ω -periodic solutions for a Stochastic fractional evolution equation driven by Levy noise with piecewise constant argument*. *arXiv : 1609.01444v1 [math.DS]*. 6 Sep 2016.
- [23] A. Lunardi. *Analytic Semigroups and Optimal Regularity in Parabolic Problems*. *PNLDE* Vol. 16, Birkhauser Verlag, Basel,1995.
- [24] J.Seidler. *Da Prato-Zzabczyk's maximal inequality revisited. I*. *Math. Bohem* 118(1), 67-106, 1993
- [25] J.Seidler and T. Sobukawa. *Exponential Integrability of Stochastic Convolutions*. *J. London Math. Soc. (2)* 67 (2003), 245-258